

SureGuard® SC

HERBICIDE



APPROVED FOR
AQUATIC &
BAREGROUND

NOW REGISTERED
IN CALIFORNIA

Residual Control that Lasts – Now in a Convenient Liquid Formula.

New from Nufarm, liquid SureGuard® SC provides an exceptional value that allows you to offer more and make more – a perfect solution for your customers. SureGuard SC delivers the same long lasting residual control of grass and broadleaf weeds that you've come to expect from SureGuard in a new convenient liquid formulation. Also, SureGuard SC enhances the speed and spectrum of glyphosate when used as a tank mix partner, even on glyphosate resistant weeds.

TOUGH CONTROL, QUICK

- **LABEL UPDATE:** Now approved for aquatic and bareground use sites
- Longest residual pre-emergent control
- Convenient easy to mix and apply patent pending liquid formulation
- Broad spectrum including tough-to-control weeds
- Fast and effective knockdown when tank mixed with glyphosate, even on glyphosate-resistant weeds
- No staining, spray formulation
- No yellow color, odor, or concerns with off-target movement and volatility

GREATER EASE, SAME POWER

USE RATES ARE EQUIVALENT TO ORIGINAL DRY FORMULATION:

1 liquid fl oz SureGuard SC = 1 dry oz SureGuard WDG

See label for complete directions for use

EPA REG. NO.	71368-114
ACTIVE INGR.	flumioxazin (44%)
FORMULATION	liquid SC
CHEM. FAMILY	N-phenylphthalimide
HRAC NO.	14
RES. USE	no
PACKAGE SIZE	8 x 1 pt, 4 x 64 fl oz case



FIELD-GROWN NURSERY



CONTAINER-GROWN NURSERY



AQUATIC



Nufarm

Grow a better tomorrow.

SureGuard® SC

HERBICIDE

KEY USES

Container and field grown conifers including Christmas trees
 Container and field grown deciduous trees and non-bearing fruit and nut trees and vines
 Non-crop areas and around buildings, fencerows, gravel surfaces, driveways, ground mats, and other similar sites
 Slow moving or quiescent aquatic sites

KEY WEEDS CONTROLLED

(See product label for complete list)

Bittercress
 Barnyardgrass*
 Bluegrass*
 Chickweed
 Crabgrass*
 Dandelion*
 Eclipta
 Field pennycress
 Florida beggarweed
 Florida pusley
 Foxtail*
 Goosegrass*
 Groundsel
 Kochia
 Lambsquarters
 Mallow
 Morningglory
 Nightshade
 Palmer amaranth
 Pennsylvania smartweed
 Pigweed
 Pineapple-weed*
 Prickly sida (*teaweed*)
 Puncturevine
 Ragweed
 Redmaids
 Russian thistle
 Shepherd's-purse
 Spiny amaranth
 Spurge
 Velvetleaf
 Waterhemp
 Yellow rocket

**Pre-emerge only*

AQUATIC WEEDS

Alligator weed
 Duckweed
 Frog's bit
 Water fern
 Water lettuce
 Watermeal

USE SUMMARY

AQUATIC AREAS

Apply this product as a broadcast spray at 6 to 12 fl oz of formulated product per acre plus an adjuvant approved for use in aquatics.

Apply this product in a minimum of 30 gallons of water per acre to all areas of the water body where weeds exist. If a second application is required to provide control, it is recommended that a treatment be made once the return of these weeds is first observed, but no sooner than 28 days after the last treatment.

Application of this product during early morning hours may enhance weed control. When applying to densely packed actively growing surface weeds, ensure adequate coverage.

This product may be tank mixed with 2,4-D, diquat, glyphosate or other registered foliar applied herbicides for enhanced control of floating and emergent weeds.

PRE-EMERGENCE APPLICATION

Pre-emergence weed control with SureGuard® SC is most effective when applied to clean, weed free soil surfaces prior to weed emergence. Moisture is necessary to activate this product on soil for residual weed control. Dry weather following application of this product may reduce effectiveness. However, when adequate moisture is received after dry conditions, this product will control susceptible germinating weeds.

To ensure uniform coverage when using boom sprayers, use 10 to 40 gallons of spray solution per acre. When making backpack applications, apply 50 to 100 gallons of spray solution per acre. Nozzle selection should meet manufacturer's gallonage and pressure recommendation for pre-emergence herbicide application.

POST-EMERGENCE APPLICATION

The most effective post-emergence weed control occurs when applied in combination with a surfactant to weeds less than 2" in height. Apply only to actively growing weeds. This product is most effective when applied under sunny conditions at temperatures above 65°F.

SureGuard SC is rainfast one hour after application. Applications should not be made if rain is expected within one hour of application or efficacy may be reduced.

To ensure thorough coverage when using boom sprayers apply 15 to 30 gallons of spray solution per acre. Apply 20 to 30 gallons per acre when using a boom sprayer if dense vegetation or heavy residue is present on the soil surface. When applying with a backpack sprayer, apply 1 gallon of spray solution per 500 to 1,000 square feet. Nozzle selection should meet manufacturer's gallonage and pressure recommendations for post-emergence herbicide application.

See label for complete use directions, restrictions and limitations.

