

SureGuard[®] SC

HERBICIDE

REDUCE LABOR AND COSTS WITH THE LONGEST RESIDUAL CONTROL.

SAVE TIME AND MONEY, GAIN FLEXIBILITY AND OPTIONS. SureGuard[®] SC Herbicide delivers the longest pre-emerge residual control of grass and broadleaf weeds – often 8 months or more – saving time and costs associated with re-application. SureGuard SC also takes down post-emerge weeds up to 2" in height for a wider application window. For powerful post-emerge knockdown with lasting residual control, SureGuard SC is compatible with non-selective herbicides like glyphosate* (Razor[®] Pro). The patent-pending liquid formulation is easy to mix, apply and clean without staining, odor, or volatility concerns. SureGuard SC is now approved for aquatic and bare ground use.

COMPETITIVE ADVANTAGES

SAVE TIME, MONEY AND LABOR

- Apply once: longest residual pre-emerge control, often 8 months or more
- Easy mix, apply and clean liquid formulation saves time
- Non-staining spray formulation with minimal color, odor, off-target movement or volatility concerns
- Broad-spectrum control including tough-to-control weeds

GAIN FLEXIBILITY AND OPTIONS

- Wider application window: unlike the competition, SureGuard SC also controls post-emerge weeds up to 2" – if timing is off, emerged weeds are still controlled
- One product for many uses: container- and field-grown ornamentals and cracks and crevices
- Mix with glyphosate* for fast and effective knockdown with longest residual, even on resistant weeds



8-MONTH CONTROL WITH 1 APPLICATION

*Do not apply glyphosate to containerized ornamentals.

AT-A-GLANCE INFO

EPA REG. NO. 71368-114

ACTIVE INGR. flumioxazin

FORMULATION liquid SC

CHEM. FAMILY N-phenylphthalimide

HRAC NO. 14

SIGNAL WORD no signal word required

RESTRICTED no

PACKAGE SIZE 8 x 1 pt, 4 x 64 fl oz

 FIELD-GROWN NURSERY

 CONTAINER-GROWN NURSERY

 PRE-EMERGE

 POST-EMERGE

 AQUATIC



SureGuard® SC

HERBICIDE

REDUCE LABOR AND COSTS WITH THE LONGEST RESIDUAL CONTROL.

KEY USES

Container and field grown conifers including Christmas trees
Container and field grown deciduous trees and non-bearing fruit and nut trees and vines
Non-crop areas and around buildings, fencerows, gravel surfaces, driveways, ground mats, and other similar sites
Slow moving or quiescent aquatic sites

KEY WEEDS CONTROLLED

(See product label for complete list)

Barnyardgrass*	Prickly sida (teaweed)
Bittercress	Puncturevine
Bluegrass	Ragweed
Chickweed	Redmaids
Crabgrass*	Russian thistle
Dandelion*	Shepherd's-purse
Eclipta	Spiny amaranth
Field pennycress	Spurge
Florida beggarweed	Velvetleaf
Florida pusley	Waterhemp
Foxtail*	Yellow rocket
Goosegrass*	AQUATIC WEEDS
Groundsel	Alligator weed
Kochia	Duckweed
Lambsquarters	Frog's bit
Morningglory	Water fern
Nightshade	Water lettuce
Palmer amaranth	Watermeal
Pigweed	*Pre-emerge only
Pineapple-weed*	

OVER-THE-TOP CONIFER TIP

SureGuard SC may be applied over-the-top to many dormant (hardened off) conifer plants. As a rule, never apply before 2 hard freezes and never after spring bud break.

32°
FREEZE 1

32°
FREEZE 2



GAIN FLEXIBILITY: A WIDER APPLICATION WINDOW



PRE-EMERGE CONTROL

Make application prior to weed emerge. Do not apply over-the-top of emerged or actively growing ornamentals. Moisture is needed to move SureGuard SC into the soil. Dry weather following application may reduce effectiveness. However, when adequate moisture is received after dry conditions, susceptible germinating weeds will be controlled.



POST-EMERGE CONTROL

For best results, apply to actively growing weeds less than 2" in height. For most effective post-emerge weed control, apply with a surfactant under sunny conditions at temperatures above 65°F. SureGuard SC is rainfast in 1 hour, do not apply if rain is expected within 1 hour.

GAIN OPTIONS: ONE PRODUCT FOR MANY USES

SureGuard SC is labeled for use on many sites. Take care of aquatics, bare grounds, landscape and container ornamentals, dormant Bermuda and more.



CONTAINER- AND FIELD-GROWN DECIDUOUS TREES AND CONIFERS

Apply as single or split applications to established trees. Do not apply to trees or transplants less than 1 year. May apply over the top only if bud break has not occurred or plants are hardened off.



BARE GROUND NON-CROP AREAS

Use for non-selective vegetation control to maintain bare ground non-crop areas. Do not apply when weather conditions favor spray drift from treated areas.



AQUATIC WEED CONTROL

Control floating or emerged weeds using surface applications or floating and submersed weeds using subsurface applications. For most effective control, apply to actively growing weeds in water with a pH of less than 8.5.



ESTABLISHED WOODY LANDSCAPE ORNAMENTALS

Apply as directed sprays around well established woody shrubs and trees. Do not apply around landscape ornamentals for at least 30 days after transplant, or at least 2 months before ornamentals will be planted.

GAIN EASE: LIQUID FORMULA FOR EASY MIX & APPLY

Convenient liquid formula makes mixing, applying and cleaning easy.



RATES ARE EQUIVALENT TO ORIGINAL DRY FORMULATION

1 liquid fl oz SureGuard SC = 1 dry oz SureGuard WDG

See label for complete directions for use.



nufarm.com/usturf

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/usturf to download a full label.

©2019 Nufarm. Important: Always read and follow label instructions. Razor® is trademark of Nufarm. SureGuard® is a trademark of Valent U.S.A. LLC.



16-GHN-0002-F